

1 **Formation of propionate and butyrate by the human colonic microbiota**

2

3

4 **Petra Louis* and Harry J. Flint**

5

6 Rowett Institute of Nutrition and Health, University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, UK

7

8 *Corresponding author; p.louis@abdn.ac.uk; phone +44 1224 438735.

9

10 Running title: Propionate and butyrate producing gut microbes

11

12 *Originality-Significance statement:* In recent years there has been a tendency to rely on sequence data to
13 assign function to microorganisms, however, this has at times led to miss-annotations and the wrong
14 conclusions being drawn. This manuscript pulls together the current knowledge on butyrate and propionate
15 metabolism in the human gut by taking account of biochemical studies performed on microorganisms in
16 addition to sequence information. In particular the areas on 1,2-propanediol metabolism and protein
17 metabolism in the gut have, to our knowledge, not been comprehensively reviewed in the context of gut
18 microbiology.

19

20 Summary

21

22 The human gut microbiota ferments dietary non-digestible carbohydrates into short-chain fatty acids
23 (SCFA). These microbial products are utilized by the host and propionate and butyrate in particular exert
24 a range of health-promoting functions. Here we provide an overview of the metabolic pathways utilized
25 by gut microbes to produce these two SCFA from dietary carbohydrates and from amino acids resulting
26 from protein breakdown. This overview emphasizes the important role played by cross-feeding of
27 intermediary metabolites (in particular lactate, succinate and 1,2-propanediol) between different gut
28 bacteria. The ecophysiology, including growth requirements and responses to environmental factors, of
29 major propionate and butyrate producing bacteria are discussed in relation to dietary modulation of
30 these metabolites. A detailed understanding of SCFA metabolism by the gut microbiota is necessary to
31 underpin effective strategies to optimize SCFA supply to the host.

32

33

34 Introduction

35

36 Short chain fatty acids (SCFA) are the major metabolic products of anaerobic fermentation by microbial
37 communities that colonize the mammalian gut, typically reaching total concentrations of 50-200 mM in the
38 human large intestine. They are taken up efficiently by the gut mucosa and have important impacts upon
39 host physiology as sources of energy, as regulators of gene expression and as signaling molecules that are
40 recognized by specific receptors (Morrison & Preston, 2016; Koh *et al.*, 2016). New mechanisms by which
41 SCFA regulate immune cell development and suppress inflammation have been uncovered recently (Louis
42 *et al.*, 2014; Richards *et al.*, 2016). It is apparent however that the three major SCFA, acetate, propionate
43 and butyrate, differ considerably in their potential effects upon host physiology. First, they differ in their
44 fate and tissue distribution, with butyrate being used preferentially as an energy source by the gut mucosa,
45 propionate contributing to gluconeogenesis in the liver and acetate achieving the highest systemic
46 concentrations in blood (Morrison & Preston, 2016). Second, there are differences in interactions with host

47 proteins (eg. inhibition of histone deacetylases by butyrate and propionate) and receptors (Bolognini *et al.*,
48 2016). This makes it particularly relevant to consider the microbial origin of these major fermentation
49 products and the potential for changes in diet and gut physiology to affect their relative production rates
50 and concentrations. This brief review will focus on butyrate and propionate as these two acids are most
51 often considered to benefit health, including protection against colorectal cancer in the case of butyrate
52 and promotion of satiety and reduction in cholesterol in the case of propionate (Morrison & Preston, 2016).
53 Acetate is a net fermentation product for most gut anaerobes that is also produced by reductive
54 acetogenesis, and almost invariably achieves the highest concentrations among the SCFA in the gut lumen.
55 In contrast, propionate and butyrate are produced by distinct subsets of gut bacteria. We consider here
56 what is currently known about the phylogenetic distribution of pathways leading to the formation of these
57 two SCFA within the human colonic microbiota and the potential for diverse dietary and environmental
58 factors to differentially modulate their production. Some fermentation products, including lactate,
59 succinate and 1,2-propanediol, do not usually accumulate to high levels in the human colon of healthy
60 adults, as they can also serve as substrates for other bacteria, including propionate and butyrate producers.
61 As the microbial metabolism of these compounds is intricately linked to the degradation of the main
62 dietary substrates, it will be discussed together with propionate and butyrate formation from
63 carbohydrates and proteins, respectively.

64

65

66 **Pathways and bacterial groups contributing to butyrate formation from carbohydrates**

67

68 Butyrate is produced from carbohydrates via glycolysis from the combination of two molecules of acetyl-
69 CoA to form acetoacetyl-CoA, followed by stepwise reduction to butyryl-CoA. Two different pathways are
70 known for the final step in butyrate formation from butyryl-CoA, which proceeds either via butyryl-
71 CoA:acetate CoA-transferase or via phosphotransbutyrylase and butyrate kinase (Louis & Flint, 2009) (Fig.
72 1). Butyrate-producing species are found interspersed with butyrate non-producing species in the two
73 predominant families of human colonic Firmicutes, *Ruminococcaceae* and *Lachnospiraceae*, as well as in

other Firmicutes families, including *Erysipelotrichaceae* and *Clostridiaceae* (Barcenilla *et al.*, 2000; Louis *et al.*, 2004). We will briefly consider the characteristics of butyrate-producers that belong to the two most abundant families of Firmicutes. We should note that, as summarized in Table 1, many dominant human colonic Firmicutes (eg. *Blautia* spp., *Eubacterium eligens*, *Ruminococcus* spp.) lack the ability to form butyrate from carbohydrates.

Ruminococcaceae. *Faecalibacterium prausnitzii*, one of the most abundant species present in the healthy human microbiota, produces butyrate via butyryl-CoA:acetate CoA-transferase with net consumption of acetate, and acetate stimulates its growth on carbohydrate energy sources (Duncan *et al.*, 2002). While *F. prausnitzii* strains are obligate anaerobes, they also show growth stimulation by low concentrations of oxygen in the presence of riboflavin and reduced compounds such as cysteine or glutathione (Khan *et al.*, 2012). It is hypothesized that this ability may provide a niche for the bacterium to thrive in the proximity of the colonic wall, where oxygen is diffusing in from the bloodstream. Oxygen consumption is accompanied by a decrease in butyrate formation (Khan *et al.*, 2012). *F. prausnitzii* isolates show limited ability to utilize dietary polysaccharides such as starch and hemicellulose for growth, but some strains utilize inulin and pectin derivatives and the ability to utilize uronic acids is widespread (Lopez-Siles *et al.*, 2012). *F. prausnitzii* is depleted in inflammatory bowel disease patients, especially Crohn's disease, and evidence that it has anti-inflammatory action has attracted interest in this species as a potential therapeutic (Quévrain *et al.*, 2016). Similarly *Butyrivibrio* *pullicaecorum* is also reported to be less abundant in inflammatory bowel disease patients, and might also have therapeutic potential (Eeckhaut *et al.*, 2013). Butyrate production has been reported for other Ruminococcaceae (Table 1), but rather little is known about most of these organisms.

Lachnospiraceae. *Eubacterium rectale* and the closely related *Roseburia* species constitute a major group of butyrate-producing Firmicutes that share the butyryl-CoA:acetate CoA-transferase route for butyrate production and the same genomic organization of their butyrate synthetic genes from acetyl-CoA to butyryl-CoA (Louis & Flint, 2009). In some *Roseburia* strains, particularly at mildly acidic pH, butyrate is almost the sole fermentation acid produced, with net consumption of acetate typically accompanying the formation of butyrate (Kettle *et al.*, 2015). Other strains and species produce formate and lactate in

101 addition to butyrate (Louis & Flint, 2009). Genome analysis reveals that there is considerable capacity
102 within this group to utilize diet-derived polysaccharides including starch, arabinoxylan and inulin, that
103 varies substantially between strains and species (Sheridan *et al.*, 2016).

104 Butyrate-producing Lachnospiraceae show considerable divergence in their phylogeny, gene
105 organization and physiology (Louis & Flint, 2009) (Table 1). Other Lachnospiraceae that possess the butyryl-
106 CoA:acetate CoA-transferase gene include *Eubacterium hallii*, *Anaerostipes hadrus*, *Coprococcus catus*, un-
107 characterised species related to isolates SS3/4 and M62/1, and some uncultured organisms (Louis *et al.*,
108 2010; Reichardt *et al.*, 2014). Two species of *Coprococcus*, in common with many *Clostridium* species that
109 belong to other families of Firmicutes, use the butyrate kinase rather than CoA-transferase enzyme for the
110 final step in butyrate formation (Louis *et al.*, 2004; Louis & Flint, 2009). *E. rectale* and *E. hallii* are among the
111 10 most abundant species reported in the human faecal microbiota (Qin *et al.*, 2010; Walker *et al.*, 2011)
112 (Table 1) and together accounted for 44% of butyryl CoA:acetate CoA-transferase sequences amplified from
113 faecal samples of 10 healthy volunteers (Louis *et al.*, 2010).

114 Lactate can be produced from carbohydrates by many different gut bacteria (Duncan *et al.*, 2004).
115 *In vitro* incubations of ¹³C lactate with human intestinal microbiota show that the label is recovered in
116 acetate, propionate and butyrate. The proportions of these products can vary widely, however, with acidic
117 pH favouring butyrate (Belenguer *et al.*, 2007). In addition, there is considerable inter-individual variation in
118 the fate of ¹³C lactate, which is assumed to reflect differences in the relative abundance of lactate-utilising
119 species within the microbiota (Bourriaud *et al.*, 2005; Morrison *et al.*, 2006). Certain Lachnospiraceae have
120 the ability to grow in the presence of lactate and acetate to produce butyrate, showing an overall net
121 stoichiometry of 4 mols of lactate and 2 mols of acetate producing 3 mols of butyrate (Duncan *et al.*, 2004).
122 These include the abundant species *A. hadrus*, which uses only D-lactate (Allen-Vercoe *et al.*, 2012) and *E.*
123 *hallii*, which is able to utilize both lactate isomers (Duncan *et al.*, 2004; Muñoz-Tamayo *et al.*, 2011). Lactate
124 oxidation to pyruvate by direct reduction of NAD⁺ is energetically unfavourable. Anaerobic lactate utilisers
125 carry a lactate dehydrogenase that operates in complex with an electron-transferring flavoprotein that
126 couples the endergonic NAD⁺ reduction to ferredoxin oxidation, in a process called electron confurcation
127 (Weghoff *et al.*, 2014).

128

129

130 **Pathways and bacterial groups contributing to propionate formation from carbohydrates**

131

132 Two pathways are known for the formation of propionate from sugar fermentation by gut bacteria. Most
133 hexose and pentose sugars are processed through the succinate pathway (Fig.2) whereas the deoxy sugars
134 fucose and rhamnose are metabolized by the propanediol pathway (Fig.3).

135 The succinate pathway is found mainly in Bacteroidetes and in the Negativicutes class of Firmicutes
136 (Reichardt *et al.*, 2014). It is the major route for propionate formation from dietary carbohydrates driven by
137 the abundant Bacteroidetes, and relative Bacteroidetes abundance was found to correlate with relative
138 faecal propionate levels in human volunteers (Salonen *et al.*, 2014). Succinate is a precursor of propionate,
139 but can accumulate in cultures of *Bacteroides* spp. under growth conditions where PEP carboxykinase is
140 repressed, eg. at high pCO₂ and high dilution rates (Caspari & Macy, 1983). Conversion of succinate to
141 propionate also requires vitamin B₁₂ and succinate has been shown to accumulate in B₁₂-depleted cultures
142 of *Prevotella ruminicola* (Strobel 1992). Some species of Bacteroidetes, notably *Prevotella copri*, apparently
143 produce succinate rather than propionate as their main fermentation product and succinate accumulation
144 has been reported particularly in the rat gut (De Vadder *et al.*, 2016). The succinate pathway is known to be
145 present in some Ruminococcaceae, such as *Ruminococcus flavefaciens*, which also produces succinate
146 rather than propionate as the end product (Macfarlane & Gibson, 1997). On the other hand, some human
147 colonic bacteria belonging to the Negativicutes class of Firmicutes (eg. *Phascolarctobacterium*
148 *succinatutens*; Watanabe *et al.*, 2012), have the ability to convert succinate to propionate (Flint *et al.*, 2014;
149 Reichardt *et al.*, 2014). This activity may explain why succinate accumulation is infrequently reported for
150 human faecal samples, although 3 of the 14 overweight human volunteers in one recent dietary study
151 showed elevated faecal succinate concentrations (>30 mM) in samples from a non-starch polysaccharide-
152 supplemented diet (reported in Salonen *et al.*, 2014, Supplementary information). Other Negativicutes
153 bacteria convert lactate to propionate either via the succinate pathway (eg. *Veillonella* spp.) or via the

154 acrylate pathway (*Megasphaera elsdenii*) (Reichardt *et al.*, 2014) (Fig. 2). The acrylate pathway has also
155 been shown to operate recently in a species of Lachnospiraceae, *Coproccoccus catus* (Reichardt *et al.*, 2014).

156

157 Formation of propionate and propanol from the deoxy sugars rhamnose and fucose via the
158 propanediol pathway has been demonstrated in dominant gut commensal bacteria belonging to the
159 Lachnospiraceae, including *Roseburia inulinivorans* and *Blautia* species (Scott *et al.*, 2006; Reichardt *et al.*,
160 2014) (Table 1, Fig. 3). Metabolism of rhamnose and fucose via this pathway has also been reported for
161 *Salmonella* and *Listeria* species (Xue *et al.*, 2008). Other bacteria, including *Bacteroides* species, *Escherichia*
162 *coli* and *Anaerostipes rhamnosivorans*, are able to degrade deoxy sugars via the propanediol pathway, but
163 produce the pathway intermediate 1,2-propanediol as the final product (Saxena *et al.*, 2010; Rodionova *et*
164 *al.*, 2013; Bui *et al.*, 2014). 1,2-propanediol can also be produced from other sugars via the glycolysis
165 intermediate dihydroxyacetone-phosphate and methylglyoxal by microbes including *Escherichia coli*,
166 *Clostridium sphenoides* and the yeast *Saccharomyces cerevisiae* (Bennett & San, 2001; Saxena *et al.*, 2010).
167 Methylglyoxal is further metabolised to 1,2-propanediol either via lactaldehyde or via hydroxyacetone (Fig.
168 3). In *C. sphenoides* it has been shown that 1,2-propanediol formation via dihydroxyacetone-phosphate
169 operates under phosphate limitation and it remains to be established whether it plays a major role in the
170 gut environment. A third pathway for 1,2-propanediol production via lactaldehyde operates from lactate in
171 *Lactobacillus buchneri*. The pathway has been elucidated in a strain isolated from maize silage (Gänzle,
172 2015), but this species has also been detected in the human gut (Mikelsaar *et al.*, 2016).

173 *E. hallii* and *Lactobacillus reuteri*, although unable to grow on fucose or rhamnose, are nevertheless
174 able to utilise 1,2-propanediol to produce propionate and propanol (Gänzle, 2015; Engels *et al.*, 2016) (Fig.
175 3). Furthermore, metagenomic mining for dehydratases has indicated that further gut anaerobes, including
176 *Flavonifractor plautii*, *Intestinimonas butyriproducing* and *Veillonella* spp. may also be able to produce
177 propionate from this substrate (Engels *et al.*, 2016). Thus, cross-feeding of the intermediate 1,2-
178 propanediol between different bacteria may play an important role in the production of propionate from
179 deoxy sugars. The conversion of 1,2-propanediol to propionate, which is dependent on vitamin B₁₂, takes
180 place in polyhedral bodies, microcompartments that sequester the toxic pathway intermediate

181 propionaldehyde (Chowdhury *et al.*, 2014). Interestingly, glycerol is converted to 1,3-propanediol and 3-
182 hydroxypropionate in *L. reuteri* and *E. hallii* by the same dehydratase that acts on 1,2-propanediol (Gänzle,
183 2015; Engels *et al.*, 2016) indicating that glycerol utilization may be the primary function of this enzyme in
184 these species. It is also worth noting that the pathway intermediate 3-hydroxypropionaldehyde, also known
185 as reuterin, is a potent antimicrobial compound (Gänzle, 2015).

186

187

188

189 **Butyrate and propionate formation from proteins and amino acids**

190

191 Propionate and butyrate are also formed as products from peptide and amino acid fermentation (Fig. 1 &
192 2), although the numbers of amino acid-fermenting bacteria have been estimated to constitute less than
193 1% of the large intestinal microbiota (Smith & Macfarlane, 1998; Dai *et al.*, 2011). It is estimated that the
194 colon receives approximately 13 g of protein and peptides per day, and large amounts of soluble protein
195 and peptides were found in intestinal contents of sudden death victims (Smith & Macfarlane, 1998).
196 Peptides seem to be preferred over free amino acids by gut bacteria. Low gut pH and the presence of
197 carbohydrates reduces peptide and amino acid fermentation *in vitro*, which helps to explain why microbial
198 amino acid fermentation is higher in the distal than the proximal colon contents (Smith & Macfarlane,
199 1998). Amino acid fermentation leads to the production of potentially harmful metabolites (for example
200 phenolic and indolic compounds, amines, ammonia) in addition to branched-chain fatty acids (BCFA) and
201 SCFA (Smith & Macfarlane, 1997; Dai *et al.*, 2011).

202 *In vitro* incubations of faecal slurries with individual amino acids showed that propionate was
203 produced mainly from aspartate, alanine, threonine and methionine, whereas butyrate was a major
204 fermentation product from glutamate, lysine, histidine, cysteine, serine and methionine (Smith &
205 Macfarlane, 1997). While several Bacteroidetes have major roles in proteolysis and in propionate formation
206 from peptides (Macfarlane & Macfarlane, 1995), certain Firmicutes species also show high activity on
207 amino acids, notably *Intestinimonas* AF211, which ferments glucose and lysine to butyrate via distinct

208 pathways (Bui *et al.*, 2015) (Fig. 1). Several different pathways exist for glutamate degradation in butyrate-
209 producing bacteria, which have mainly been studied in *Clostridium* species not originating from gut
210 environments (Barker, 1981; Buckel, 2001). However, there is genomic and metagenomic evidence that
211 they are also present in some gut bacteria (Potrykus *et al.*, 2008; Vital *et al.*, 2014). The glutamate
212 degradation pathways enter the main butyrate pathway either via pyruvate (3-methylaspartate pathway;
213 *Clostridium limosum*, *Fusobacterium* spp.) or crotonyl-CoA (4-aminobutyrate pathway, discussed in more
214 detail below, and 2-hydroxyglutarate pathway, found in different Firmicutes including *Acidaminococcus*
215 *fermentans*, *Clostridium sporosphaeroides*, *Clostridium symbiosum*, *Fusobacterium* spp. and
216 *Peptostreptococcus asaccharolyticus*) (Fig. 1). Some bacteria belonging to the Acidaminococcaceae also
217 degrade glutamate via the 3-methylaspartate pathway, but produce propionate rather than butyrate from
218 the intermediate pyruvate (Buckel, 2001) (Fig. 2).

219 Glutamate degradation to 4-aminobutyrate (gamma-aminobutyrate, GABA) is carried out under
220 acid stress to maintain intracellular pH homeostasis in a number of gut bacteria (Feehily & Karatzas, 2013),
221 and a bacterial isolate exclusively growing on GABA has recently been found
222 (<http://www.abstractsonline.com/pp8/#!/4060/presentation/18619>). As GABA also acts as a
223 neurotransmitter, the abundance of microbes involved in the production or consumption of GABA may
224 influence mood and behaviour. The pathway for GABA degradation is shared with succinate degradation via
225 succinate semialdehyde and 4-hydroxybutyrate (Fig. 1), and butyrate production from succinate via this
226 pathway has been demonstrated in *Porphyromonas gingivalis* and *Clostridioides difficile* (Ferreyra *et al.*,
227 2014; Yoshida *et al.*, 2016).

228 The fermentation routes of other amino acids are less well understood. Histidine is converted to
229 glutamate (Potrykus *et al.*, 2008; Kanehisa *et al.*, 2016), which is in agreement with high levels of butyrate
230 being formed from histidine by faecal microbiota (Smith & Macfarlane, 1997). Alanine, serine and cysteine
231 are broken down to pyruvate (Potrykus *et al.*, 2008; Carbonero *et al.*, 2012), thus product formation
232 depends on the bacterium utilizing those amino acids and their corresponding fermentative pathways. For
233 example, in *Clostridium propionicum*, alanine fermentation leads to the production of propionate via
234 pyruvate, lactate and the acrylate pathway (Buckel, 2001) (Fig. 2). Threonine and methionine are converted

235 to 2-oxobutyrate, which leads to propionate formation (Fig. 2) (Barker, 1981; Smith & Macfarlane, 1997;
236 Kanehisa *et al.*, 2016). Several routes for the breakdown of aspartate exist, via alanine, threonine,
237 oxaloacetate or fumarate (Smith & Macfarlane, 1997; Kanehisa *et al.*, 2016) (Fig. 2), which accounts for the
238 fact that it is mainly converted to propionate in *in vitro* incubations.

239

240

241 **Role of CoA-transferases in SCFA metabolism**

242

243 Propionate and butyrate can be generated from their respective CoA thioesters either by transfer of the
244 CoA-moiety onto another metabolite, or by conversion via propionyl-phosphate or butyryl-phosphate. The
245 second (kinase) route leads to the generation of ATP, but the CoA-transferase route also conserves the
246 energy of the CoA bond in the newly formed CoA-derivative of the co-substrate. Acetate is a common co-
247 substrate in CoA-transferase reactions, and the high acetate concentrations in the large intestine provide a
248 possible explanation for the prevalence of the butyryl-CoA:acetate CoA-transferase route in gut microbes
249 (Louis *et al.*, 2004) (see also section on pH below). Bacteria often carry multiple different CoA-transferases
250 in their genomes, with *Intestinimonas* AF211 encoding at least 14 such enzymes (Bui *et al.*, 2015). It can be
251 difficult to pin-point which gene is responsible for SCFA formation, especially as CoA-transferases tend to
252 have broad substrate specificity. For example, the purified butyryl-CoA:acetate CoA-transferase (*butCoAT*
253 gene product) from *Roseburia hominis* has a similar affinity for butyryl-CoA and propionyl-CoA although the
254 enzyme is clearly responsible for butyrate formation in this species (Charrier *et al.*, 2006) (Table 2). Gene
255 expression evidence in *Intestinimonas* AF211 suggested that the enzyme AtoD-A, responsible for butyryl
256 CoA:acetoacetate CoA-transferase activity, plays a key role in conversion of lysine to butyrate, while the
257 ButCoA gene product mediated the final step in butyrate formation from glucose (Bui *et al.*, 2015). In
258 *Clostridium aminobutyricum*, a CoA-transferase that acts on 4-hydroxybutyrate and butyryl-CoA links the
259 final step of butyrate production to the formation of 4-hydroxybutyryl-CoA further up in the glutamate
260 fermentation pathway (Buckel, 2001). Similarly, *C. propionicum* links the formation of lactoyl-CoA in the
261 acrylate pathway to propionate formation via a CoA-transferase (Buckel, 2001). There are also instances

262 where different CoA-transferases appear to have evolved for the same enzymatic reaction. Thus, bacteria
263 belonging to the Erysipelotrichaceae do not carry a gene closely related to the butyryl-CoA:acetate CoA-
264 transferase identified in other Firmicutes. Instead, a gene more closely related to propionate CoA-
265 transferases is thought to be responsible for butyrate formation in these organisms (Eeckhaut *et al.*, 2011).

266

267

268 **Impact of the gut environment**

269

270 **pH.** Gut pH has a major impact on competition between different groups of bacteria within the microbial
271 community. In pH-controlled *in vitro* continuous culture experiments with soluble polysaccharide provided
272 as the main energy source, mildly acidic pH has been shown to curtail the growth of *Bacteroides* spp.
273 relative to Firmicutes and Actinobacteria (Walker *et al.*, 2005; Chung *et al.*, 2016). This is because human
274 colonic *Bacteroides* spp. are generally less able than many dominant Firmicutes to tolerate the presence of
275 short chain fatty acids at pH 5.5 (Duncan *et al.*, 2009). This selective inhibition and the resulting shift in
276 community composition has the consequence of limiting propionate formation and enhancing butyrate
277 production by the community at pH values around 5.5 compared with 6.5-6.8 (Walker *et al.*, 2005; Chung *et*
278 *al.*, 2016). The impact of pH shifts upon experimentally observed butyrate and propionate concentrations
279 has been successfully modelled mathematically, based on the differing tolerance to low pH of the major
280 bacterial functional groups that comprise the human colonic microbiota (Kettle *et al.*, 2015).

281 For bacteria that use the butyryl-CoA:acetate CoA-transferase route, acetate consumption and
282 butyrate production are reported to increase at mildly acidic pH compared with near neutral pH (Kettle *et*
283 *al.*, 2015). Although conversion of glucose to butyrate, 2 CO₂ and 2 H₂ can occur with no net uptake of
284 acetate (Gottschalk, 1979), net acetate uptake is typically observed for species of *Roseburia* and *F.*
285 *prausnitzii*. Theoretical stoichiometries involving net acetate uptake are shown in Fig. 4A, which also
286 assumes that some of the reducing power that is generated drives proton export, increasing the ATP yield
287 per glucose fermented (Buckel & Thauer, 2013). Incorporation of exogenous acetate via the CoA-
288 transferase reaction results in some loss of ATP production via acetyl-phosphate, but this is more than

289 compensated by the additional ATP formed from proton export, giving a potential maximum of 4 ATP
290 formed per glucose metabolized when 2 mols of acetate are taken in for each mol of glucose fermented.
291 Interestingly, Fig. 4B shows that the predictions from these stoichiometries (based on the generalised
292 equation shown in Fig. 4A) fit experimental data for the impact of pH on metabolites produced by *F.*
293 *prausnitzii* and two *Roseburia* spp. in anaerobic batch culture (Kettle *et al.*, 2015). Thus low pH (5.5) tends
294 to increase acetate uptake and butyrate production while near neutral pH (6.7) has the opposite effect. It
295 seems possible that the increased ATP gain associated with net acetate uptake helps to compensate for the
296 effects of low pH and might account for the reliance in the CoA-transferase route for butyrate formation in
297 these bacteria.

298 **Growth requirements.** It has been show in a rodent model that limitation of dietary iron intake can
299 dramatically decrease the production of both butyrate and propionate as lactobacilli and Proteobacteria
300 are favoured (Dostal *et al.*, 2012). Populations of *Roseburia*-related butyrate producers appear particularly
301 sensitive to iron availability, while in pure cultures of *R. intestinalis* butyrate production was favoured at
302 high iron concentrations with a switch to lactate production under iron-deficient conditions (Dostal *et al.*,
303 2015). It remains to be established whether other growth factors also have a major impact on SCFA
304 formation.

305 **Intestinal gases.** SCFA formation is also likely to be affected by differences in oxygen concentration
306 in different regions and micro-compartments of the gut due to differences in oxygen sensitivity and
307 metabolic capacity between microbes, as exemplified by the peculiar relationship of *F. prausnitzii* with
308 oxygen (discussed above). Furthermore, the abundance of microbes consuming hydrogen and thereby
309 influencing the hydrogen partial pressure in the gut also influences SCFA formation, as this affects the
310 overall balance of fermentation products formed (Macfarlane & Macfarlane, 2003; Wolf *et al.*, 2016).

311

312

313 **Concluding remarks**

314

315 Huge advances have been made in recent years in our understanding of SCFA metabolism in the human
316 gut, and many of the dominant propionate- and butyrate-producing bacteria are available in culture,
317 enabling detailed investigations into their metabolism. Recent work has emphasized that butyrate and
318 propionate can arise from fermentation both of amino acids and of carbohydrates, but the relative
319 contributions of protein and carbohydrate fermentation *in vivo* over the wide range of 'normal' human
320 dietary intakes is not yet clear. We know that high protein, low carbohydrate weight loss diets lead to a
321 disproportionate decrease in butyrate among total faecal SCFA, together with an increased proportion of
322 branched chain fatty acids that are wholly derived from branched chain amino acids and therefore provide
323 an indicator of protein fermentation (Duncan *et al.*, 2007, Russell *et al.*, 2011). This suggests strongly that
324 butyrate production is mainly determined by the supply of non-digestible carbohydrates, rather than by
325 protein fermentation. This may however reflect the particular ecology of butyrate-producing bacteria, as
326 discussed above. In the case of propionate, on the other hand, the major producers of propionate from
327 dietary carbohydrates, the Bacteroidetes, are also important peptide fermenters and the propionate
328 proportion among faecal SCFA was not decreased by such low carbohydrate diets (Duncan *et al.*, 2007). It is
329 also clear that compounds normally regarded as intermediates (eg. succinate, lactate) may accumulate in
330 certain individuals or in particular conditions. This makes it important also to consider the impacts of these
331 metabolites on the host, as for example in the case of succinate which it is suggested may provide health
332 benefits (De Vadder *et al.*, 2016). Lactate is detected as a major fermentation product in breast-fed infants
333 whose microbiota is dominated by *Bifidobacterium* spp. In adults, however, lactate accumulation is
334 associated with dysbiosis, eg, in severe colitis (Hove *et al.*, 1994), that may result in part from a lack of
335 lactate-utilizing bacteria (Belenguer *et al.*, 2007).

336 The ever-increasing availability of genomic and metagenomics sequences is a highly useful resource
337 to foster our understanding of microbial metabolism in the gut, but care has to be taken with assigning
338 function to genes by sequence analysis, which should ideally be complemented by evidence from genetic or
339 enzymatic studies. A renewed interest in isolation and study of gut bacteria (Walker *et al.*, 2014, Browne *et*
340 *al.*, 2016) together with novel systems for gene transfer and knockout on the horizon will enable a
341 thorough understanding of the different members of the microbial community. This will benefit *in vitro* and

342 *in vivo* microbial community-based studies to foster our understanding of the different ecological niches of
343 the community members, how they interact with each other and how we can modulate the system by
344 dietary means to optimize SCFA production. The fact that, in general, different phylogenetic groups of
345 bacteria are responsible for butyrate and propionate production suggests that there may be scope for
346 differentially manipulating their production by the gut microbiota.

347

348

349 **Acknowledgements**

350 The authors receive financial support from the Scottish Government Rural and Environmental Sciences and
351 Analytical Services (RESAS). We would like to thank Sylvia Duncan for SCFA data of *F. prausnitzii* grown
352 under different pH regimes and for critically reading the manuscript.

353

354 The authors declare that they have no conflicts of interest.

355

356

357 **References**

358

359 Allen-Vercoe, E., Daigneault, M., White, A., Panaccione, R., Duncan, S. H., Flint, H. J. et al. (2012)

360 *Anaerostipes hadrus* comb. nov., a dominant species within the human colonic microbiota;

361 reclassification of *Eubacterium hadrum* Moore et al. 1976. *Anaerobe* **18**: 523-529.

362 Barcenilla, A., Pryde, S. E., Martin, J. C., Duncan, S. H., Stewart, C. S., Henderson, C., & Flint, H. J. (2000)

363 Phylogenetic relationships of butyrate-producing bacteria from the human gut. *Appl Environ Microbiol*

364 **66**: 1654-1661.

365 Barker, H. A. (1981) Amino acid degradation by anaerobic bacteria. *Annu Rev Biochem* **50**: 23-40.

366 Belenguer, A., Duncan, S. H., Holtrop, G., Anderson, S. E., Lobley, G. E., & Flint, H. J. (2007) Impact of pH on
 367 lactate formation and utilization by human fecal microbial communities. *Appl Environ Microbiol* **73**:
 368 6526-6533.

369 Bennett, G. N., & San, K. Y. (2001) Microbial formation, biotechnological production and applications of 1,2-
 370 propanediol. *Appl Microbiol Biotechnol* **55**: 1-9.

371 Bolognini, D., Tobin, A. B., Milligan, G., & Moss, C. E. (2016) The pharmacology and function of receptors for
 372 short-chain fatty acids. *Mol Pharmacol* **89**: 388-398.

373 Bourriaud, C., Robins, R. J., Martin, L., Kozlowski, F., Tenailleau, E., Cherbut, C., & Michel, C. (2005) Lactate
 374 is mainly fermented to butyrate by human intestinal microfloras but inter-individual variation is
 375 evident. *J Appl Microbiol* **99**: 201-212.

376 Browne, H. P., Forster, S. C., Anonye, B. O., Kumar, N., Neville, B. A., Stares, M. D., Goulding, D., & Lawley, T.
 377 D. (2016) Culturing of 'unculturable' human microbiota reveals novel taxa and extensive sporulation.
 378 *Nature* **533**: 543-546.

379 Buckel, W. (2001) Unusual enzymes involved in five pathways of glutamate fermentation. *Appl Microbiol*
 380 *Biotechnol* **57**: 263-273.

381 Buckel, W., & Thauer, R. K. (2013) Energy conservation via electron bifurcating ferredoxin reduction and
 382 proton/Na⁺ translocating ferredoxin oxidation. *BBA-Bioenergetics* **1827**: 94-113.

383 Bui, T. P. N., de Vos, W. M., & Plugge, C. M. (2014) *Anaerostipes rhamnosivorans* sp. nov., a human
 384 intestinal, butyrate-forming bacterium. *Int J Syst Evol Microbiol* **64**: 787-793.

385 Bui, T. P. N., Ritari, J., Boeren, S., De Waard, P., Plugge, C. M., & De Vos, W. M. (2015) Production of
 386 butyrate from lysine and the Amadori product fructoselysine by a human gut commensal. *Nat*
 387 *Commun* **6**: 10062.

388 Carbonero, F., Benefiel, A. C., Alizadeh-Ghamsari, A. H., & Gaskins, H. R. (2012) Microbial pathways in
389 colonic sulfur metabolism and links with health and disease. *Front Physiol* **3**: 448.

390 Caspari, D., & Macy, J. M. (1983) The role of carbon dioxide in glucose metabolism of *Bacteroides fragilis*.
391 *Arch Microbiol* **135**: 16-24.

392 Charrier, C., Duncan, G. J., Reid, M. D., Rucklidge, G. J., Henderson, D., Young, P. et al. (2006) A novel class
393 of CoA-transferase involved in short-chain fatty acid metabolism in butyrate-producing human colonic
394 bacteria. *Microbiology* **152**: 179-185.

395 Chowdhury, C., Sinha, S., Chun, S., Yeates, T. O., & Bobik, T. A. (2014) Diverse bacterial microcompartment
396 organelles. *Microbiol Mol Biol Rev* **78**: 438-468.

397 Chung, W. S. F., Walker, A. W., Louis, P., Parkhill, J., Vermeiren, J., Bosscher, D. et al. (2016) Modulation of
398 the human gut microbiota by dietary fibres occurs at the species level. *BMC Biol* **14**: 3.

399 Dai, Z. L., Wu, G., & Zhu, W. Y. (2011) Amino acid metabolism in intestinal bacteria: Links between gut
400 ecology and host health. *Front Biosci* **16**: 1768-1786.

401 De Maesschalck, C., Van Immerseel, F., Eeckhaut, V., De Baere, S. D., Cnockaert, M., Croubels, S. et al.
402 (2014) *Faecalicoccus acidiformans* gen. nov., sp. nov., Isolated from the chicken caecum, and
403 reclassification of *Streptococcus pleomorphus* (Barnes et al. 1977), *Eubacterium biforme* (Eggerth
404 1935) and *Eubacterium cylindroides* (Cato et al. 1974) as *Faecalicoccus pleomorphus* comb. nov.,
405 *Holdemanella biformis* gen. nov., Comb. nov. and *Faecalitalea cylindroides* gen. nov., Comb. nov.,
406 respectively, within the family *Erysipelotrichaceae*. *Int J Syst Evol Microbiol* **64**: 3877-3884.

407 De Vadder, F., Kovatcheva-Datchary, P., Zitoun, C., Duchampt, A., Bäckhed, F., & Mithieux, G. (2016)
408 Microbiota-produced succinate improves glucose homeostasis via intestinal gluconeogenesis. *Cell*
409 *Metabolism* **24**: 151-157.

410 Dostal, A., Lacroix, C., Bircher, L., Pham, V. T., Follador, R., Zimmermann, M. B., & Chassard, C. (2015) Iron
 411 modulates butyrate production by a child gut microbiota in vitro. *mBio* **6**: e01453-15.

412 Dostal, A., Chassard, C., Hilty, F. M., Zimmermann, M. B., Jaeggi, T., Rossi, S., & Lacroix, C. (2012) Iron
 413 depletion and repletion with ferrous sulfate or electrolytic iron modifies the composition and
 414 metabolic activity of the gut microbiota in rats. *J Nutr* **142**: 271-277.

415 Duncan, S. H., Belenguer, A., Holtrop, G., Johnstone, A. M., Flint, H. J., & Lobley, G. E. (2007) Reduced
 416 dietary intake of carbohydrates by obese subjects results in decreased concentrations of butyrate and
 417 butyrate-producing bacteria in feces. *Appl Environ Microbiol* **73**: 1073-1078.

418 Duncan, S. H., Louis, P., & Flint, H. J. (2004) Lactate-utilizing bacteria, isolated from human feces, that
 419 produce butyrate as a major fermentation product. *Appl Environ Microbiol* **70**: 5810-5817.

420 Duncan, S. H., Louis, P., Thomson, J. M., & Flint, H. J. (2009) The role of pH in determining the species
 421 composition of the human colonic microbiota. *Environ Microbiol* **11**: 2112-2122.

422 Duncan, S. H., Hold, G. L., Harmsen, H. J. M., Stewart, C. S., & Flint, H. J. (2002) Growth requirements and
 423 fermentation products of *Fusobacterium prausnitzii*, and a proposal to reclassify it as *Faecalibacterium*
 424 *prausnitzii* gen. nov., comb. nov. *Int J Syst Evol Microbiol* **52**: 2141-2146.

425 Eeckhaut, V., van Immerseel, F., Croubels, S., de Baere, S., Haesebrouck, F., Ducatelle, R. et al. (2011)
 426 Butyrate production in phylogenetically diverse Firmicutes isolated from the chicken caecum.
 427 *Microbial Biotechnol* **4**: 503-512.

428 Eeckhaut, V., Machiels, K., Perrier, C., Romero, C., Maes, S., Flahou, B. et al. (2013) *Butyricicoccus*
 429 *pullicaecorum* in inflammatory bowel disease. *Gut* **62**: 1745-1752.

430 Engels, C., Ruscheweyh, H. J., Beerenwinkel, N., Lacroix, C., & Schwab, C. (2016) The common gut microbe
 431 *Eubacterium hallii* also contributes to intestinal propionate formation. *Front Microbiol* **7**: 713.

432 Feehily, C., & Karatzas, K. A. G. (2013) Role of glutamate metabolism in bacterial responses towards acid
 433 and other stresses. *J Appl Microbiol* **114**: 11-24.

434 Ferreyra, J. A., Wu, K. J., Hryckowian, A. J., Bouley, D. M., Weimer, B. C., & Sonnenburg, J. L. (2014) Gut
 435 microbiota-produced succinate promotes *C. difficile* infection after antibiotic treatment or motility
 436 disturbance. *Cell Host Microbe* **16**: 770-777.

437 Flint, H. J., Duncan, S. H., Scott, K. P., & Louis, P. (2014) Links between diet, gut microbiota composition and
 438 gut metabolism. *Proc Nutr Soc* **74**: 13-22.

439 Gänzle, M. G. (2015) Lactic metabolism revisited: Metabolism of lactic acid bacteria in food fermentations
 440 and food spoilage. *Curr Opin Food Sci* **2**: 106-117.

441 Gottschalk, G. (1979) *Bacterial metabolism*. Springer Verlag, New York, Heidelberg, Berlin.

442 Hove, H., Nordgaard Andersen, I., & Mortensen, P. B. (1994) Fecal DL-lactate concentration in 100
 443 gastrointestinal patients. *Scand J Gastroenterol* **29**: 255-259.

444 Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., & Tanabe, M. (2016) KEGG as a reference resource
 445 for gene and protein annotation. *Nucleic Acids Res* **44**: D457-D462.

446 Kettle, H., Louis, P., Holtrop, G., Duncan, S. H., & Flint, H. J. (2015) Modelling the emergent dynamics and
 447 major metabolites of the human colonic microbiota. *Environ Microbiol* **17**: 1615-1630.

448 Khan, M. T., Duncan, S. H., Stams, A. J. M., van Dijk, J. M., Flint, H. J., & Harmsen, H. J. M. (2012) The gut
 449 anaerobe *Faecalibacterium prausnitzii* uses an extracellular electron shuttle to grow at oxic-anoxic
 450 interphases. *ISME J* **6**: 1578-1585.

451 Koh, A., De Vadder, F., Kovatcheva-Datchary, P., & Bäckhed, F. (2016) From dietary fiber to host physiology:
 452 short-chain fatty acids as key bacterial metabolites. *Cell* **165**: 1332-1345.

453 Lopez-Siles, M., Khan, T. M., Duncan, S. H., Harmsen, H. J. M., Garcia-Gil, L. J., & Flint, H. J. (2012) Cultured
 454 representatives of two major phylogroups of human colonic *Faecalibacterium prausnitzii* can utilize
 455 pectin, uronic acids, and host-derived substrates for growth. *Appl Environ Microbiol* **78**: 420-428.

456 Louis, P., & Flint, H. J. (2009) Diversity, metabolism and microbial ecology of butyrate-producing bacteria
 457 from the human large intestine. *FEMS Microbiol Lett* **294**: 1-8.

458 Louis, P., Hold, G. L., & Flint, H. J. (2014) The gut microbiota, bacterial metabolites and colorectal cancer.
 459 *Nat Rev Microbiol* **12**: 661-672.

460 Louis, P., Young, P., Holtrop, G., & Flint, H. J. (2010) Diversity of human colonic butyrate-producing bacteria
 461 revealed by analysis of the butyryl-CoA:acetate CoA-transferase gene. *Environ Microbiol* **12**: 304-314.

462 Louis, P., Duncan, S. H., McCrae, S. I., Millar, J., Jackson, M. S., & Flint, H. J. (2004) Restricted distribution of
 463 the butyrate kinase pathway among butyrate-producing bacteria from the human colon. *J Bacteriol*
 464 **186**: 2099-2106.

465 Macfarlane, G. T., & Gibson, G. R. (1997) Carbohydrate fermentation, energy transduction and gas
 466 metabolism in the human large intestine. In *Gastrointestinal microbiology vol. 1*. R. I. Mackie, & B. A.
 467 White (eds). London, Chapman and Hall, pp. 269-318.

468 Macfarlane, S., & Macfarlane, G. T. (2003) Session: Short-chain fatty acids. Regulation of short-chain fatty
 469 acid production. *Proc Nutr Soc* **62**: 67-72.

470 Macfarlane, S., & Macfarlane, G. T. (1995) Proteolysis and amino acid fermentation. In *Human colonic*
 471 *bacteria - role in nutrition, physiology and pathology*. G. R. Gibson, & G. T. Macfarlane (eds). CRC Press
 472 Inc, pp. 75-100.

473 Mikelsaar, M., Sepp, E., Štšepetova, J., Songisepp, E., & Mändar, R. (2016) Biodiversity of intestinal lactic
 474 acid bacteria in the healthy population. *Adv Exp Med Biol* [Epublication ahead of print]

475 Morrison, D. J., & Preston, T. (2016) Formation of short chain fatty acids by the gut microbiota and their
 476 impact on human metabolism. *Gut Microbes* **7**: 189-200.

477 Morrison, D. J., Mackay, W. G., Edwards, C. A., Preston, T., Dodson, B., & Weaver, L. T. (2006) Butyrate
 478 production from oligofructose fermentation by the human faecal flora: What is the contribution of
 479 extracellular acetate and lactate? *Br J Nutr* **96**: 570-577.

480 Muñoz-Tamayo, R., Laroche, B., Walter, É., Doré, J., Duncan, S. H., Flint, H. J., & Leclerc, M. (2011) Kinetic
 481 modelling of lactate utilization and butyrate production by key human colonic bacterial species. *FEMS*
 482 *Microbiol Ecol* **76**: 615-624.

483 Potrykus, J., White, R. L., & Bearne, S. L. (2008) Proteomic investigation of amino acid catabolism in the
 484 indigenous gut anaerobe *Fusobacterium varium*. *Proteomics* **8**: 2691-2703.

485 Qin, J., Li, R., Raes, J., Arumugam, M., Burgdorf, K. S., Manichanh, C. et al. (2010) A human gut microbial
 486 gene catalogue established by metagenomic sequencing. *Nature* **464**: 59-65.

487 Quévrain, E., Maubert, M. A., Michon, C., Chain, F., Marquant, R., Tailhades, J. et al. (2016) Identification of
 488 an anti-inflammatory protein from *Faecalibacterium prausnitzii*, a commensal bacterium deficient in
 489 Crohn's disease. *Gut* **65**: 415-425.

490 Reichardt, N., Duncan, S. H., Young, P., Belenguer, A., McWilliam Leitch, C., Scott, K. P. et al. (2014)
 491 Phylogenetic distribution of three pathways for propionate production within the human gut
 492 microbiota. *ISME J* **8**: 1323-1335.

493 Richards, J. L., Yap, Y. A., McLeod, K. H., MacKay, C. R., & Marinó, E. (2016) Dietary metabolites and the gut
 494 microbiota: An alternative approach to control inflammatory and autoimmune diseases. *Clin Transl*
 495 *Immunology* **5**: e82.

496 Rodionova, I. A., Li, X., Thiel, V., Stolyar, S., Stanton, K., Fredrickson, J. K. et al. (2013) Comparative genomics
 497 and functional analysis of rhamnose catabolic pathways and regulons in bacteria. *Front Microbiol* **4**:
 498 407.

499 Russell, W. R., Gratz, S. W., Duncan, S. H., Holtrop, G., Ince, J., Scobbie, L., Duncan, G., Johnstone, A. M.,
 500 Lobley, G. E., Wallace, R. J., Duthie, G. G., & Flint, H. J. (2011) High-protein, reduced-carbohydrate
 501 weight-loss diets promote metabolite profiles likely to be detrimental to colonic health. *Am J Clin Nutr*
 502 **93**: 1062-1072.

503 Salonen, A., Lahti, L., Salojärvi, J., Holtrop, G., Korpela, K., Duncan, S. H. et al. (2014) Impact of diet and
 504 individual variation on intestinal microbiota composition and fermentation products in obese men.
 505 *ISME J* **8**: 2218-2230.

506 Saxena, R. K., Anand, P., Saran, S., Isar, J., & Agarwal, L. (2010) Microbial production and applications of 1,2-
 507 propanediol. *Indian J Microbiol* **50**: 2-11.

508 Scott, K. P., Martin, J. C., Campbell, G., Mayer, C. D., & Flint, H. J. (2006) Whole-genome transcription
 509 profiling reveals genes up-regulated by growth on fucose in the human gut bacterium "*Roseburia*
 510 *inulinivorans*". *J Bacteriol* **188**: 4340-4349.

511 Sheridan, P. O., Martin, J. C., Lawley, T. D., Browne, H. P., Harris, H. M. B., Bernalier-Donadille, A. et al.
 512 (2016) Polysaccharide utilization loci and nutritional specialization in a dominant group of butyrate-
 513 producing human colonic *Firmicutes*. *Microbial Genomics* **2**: 000043.

514 Smith, E. A., & Macfarlane, G. T. (1998) Enumeration of amino acid fermenting bacteria in the human large
 515 intestine: Effects of pH and starch on peptide metabolism and dissimilation of amino acids. *FEMS*
 516 *Microbiol Ecol* **25**: 355-368.

517 Smith, E. A., & Macfarlane, G. T. (1997) Dissimilatory amino acid metabolism in human colonic bacteria.
 518 *Anaerobe* **3**: 327-337.

519 Strobel, H. J. (1992) Vitamin B₁₂-dependent propionate production by the ruminal bacterium *Prevotella*
520 *ruminicola* 23. *Appl Environ Microbiol* **58**: 2331-2333.

521 Vital, M., Howe, A. C., & Tiedje, J. M. (2014) Revealing the bacterial butyrate synthesis pathways by
522 analyzing (meta)genomic data. *mBio* **5**: e00889.

523 Walker, A. W., Duncan, S. H., Carol McWilliam Leitch, E., Child, M. W., & Flint, H. J. (2005) pH and peptide
524 supply can radically alter bacterial populations and short-chain fatty acid ratios within microbial
525 communities from the human colon. *Appl Environ Microbiol* **71**: 3692-3700.

526 Walker, A. W., Ince, J., Duncan, S. H., Webster, L. M., Holtrop, G., Ze, X. et al. (2011) Dominant and diet-
527 responsive groups of bacteria within the human colonic microbiota. *ISME J* **5**: 220-230.

528 Walker, A. W., Duncan, S. H., Louis, P., & Flint, H. J. (2014) Phylogeny, culturing, and metagenomics of the
529 human gut microbiota. *Trends Microbiol* **22**: 267-274.

530 Watanabe, Y., Nagai, F., & Morotomi, M. (2012) Characterization of *Phascolarctobacterium succinatutens*
531 sp. Nov., an asaccharolytic, succinate-utilizing bacterium from human feces. *Appl Environ Microbiol* **78**:
532 511-518.

533 Weghoff, M. C., Bertsch, J., & Müller, V. (2014) A novel mode of lactate metabolism in strictly anaerobic
534 bacteria. *Environ Microbiol* **17**: 670-677.

535 Wolf, P. G., Biswas, A., Morales, S. E., Greening, C., & Gaskins, H. R. (2016) H₂ metabolism is widespread and
536 diverse among human colonic microbes. *Gut Microbes* **7**: 235-245.

537 Xue, J., Murrieta, C. M., Rule, D. C., & Miller, K. W. (2008) Exogenous or L-rhamnose-derived 1,2-
538 propanediol is metabolized via a *pduD*-dependent pathway in *Listeria innocua*. *Appl Environ Microbiol*
539 **74**: 7073-7079.

540 Yoshida, Y., Sato, M., Kezuka, Y., Hasegawa, Y., Nagano, K., Takebe, J., & Yoshimura, F. (2016) Acyl-CoA
541 reductase PGN-0723 utilizes succinyl-CoA to generate succinate semialdehyde in a butyrate-producing
542 pathway of *Porphyromonas gingivalis*. *Arch Biochem Biophys* **596**: 138-148.

543 Zhernakova, A., Kurilshikov, A., Bonder, M. J., Tigchelaar, E. F., Schirmer, M., Vatanen, T. et al. (2016)
544 Population-based metagenomics analysis reveals markers for gut microbiome composition and
545 diversity. *Science* **352**: 565-569.

546 Figure legends

547

548 **Fig. 1.** Microbial pathways for butyrate formation from carbohydrates, organic acids, glutamate and lysine
549 in gut communities. Carbohydrate fermentation to pyruvate via glycolysis is shown in green, butyrate
550 formation from acetyl-CoA in black, amino acid fermentation pathways in blue (intermediates after which
551 the different glutamate pathways are named are highlighted), and lactate and succinate fermentation in
552 purple and pink, respectively. See main text for key enzymes and bacteria harbouring the different
553 pathways. Redox reactions which involve electron carriers are indicated by [H]. CoA-transferase-mediated
554 reactions are indicated by ●. As indicated, co-substrates other than acetate may operate in CoA-transferase
555 reactions in some bacteria (for further detail see main text). CoA, coenzyme A; P, bound phosphate; Pi,
556 inorganic phosphate; PEP, phosphoenolpyruvate; (B₁₂), enzyme dependent on vitamin B₁₂. Dotted line
557 indicates that several intermediate steps are involved.

558

559 **Fig. 2.** Microbial pathways for propionate formation from carbohydrates, organic acids and amino acids. As
560 indicated, amino acids capable of conversion to pyruvate can also give rise to butyrate (Fig. 1).
561 Carbohydrate fermentation to pyruvate via glycolysis is shown in green, propionate formation via the
562 succinate pathway in black, amino acid fermentation pathways in blue, and acrylate pathway for lactate
563 utilisation in purple. See main text for key enzymes and bacteria harbouring the different pathways. Redox
564 reactions which involve electron carriers are indicated by [H]. CoA-transferase-mediated reactions are
565 indicated by ● (*may be performed by a CoA-transferase or CoA-ligase reaction). Propionate formation

566 from propionyl-CoA in the succinate pathway may involve either a CoA-transferase or phosphate
567 propanoyltransferase/propionate kinase reaction. CoA, coenzyme A; PEP, phosphoenolpyruvate; (B₁₂),
568 dependent on vitamin B₁₂. Dotted lines indicate that several intermediate steps are involved.

569

570 **Fig. 3.** Microbial pathways for propionate formation via 1,2-propanediol. Carriage of the different pathways
571 in gut microbes is indicated by colour. Redox reactions which involve electron carriers are indicated by [H].
572 Propionate formation from propionyl-CoA may involve either a CoA-transferase or phosphate
573 propanoyltransferase/propionate kinase reaction (indicated by a dashed line). Grey hexagon indicates that
574 the reaction is carried out in polyhedral bodies to sequester toxic intermediate propionaldehyde. CoA,
575 coenzyme A; P, bound phosphate; (B₁₂), dependent on vitamin B₁₂. Dotted line indicates that several
576 intermediate steps are involved.

577

578 **Fig. 4.** Butyrate production in bacteria that use the butyryl-CoA:acetate CoA-transferase route. A: General
579 equation for the relationship between acetate consumption and butyrate production, assuming no lactate
580 or formate are produced (modified from Louis & Flint 2009 and Kettle et al. 2015). Etf, electron-transferring
581 flavoprotein; Fd, ferredoxin; P, bound phosphate; Pi, inorganic phosphate. B: Alternative stoichiometries
582 for butyrate production based on A. Experimental data (coloured symbols) refer to *R. intestinalis* L1-82, *R.*
583 *hominis* A2-183 and *F. prausnitzii* A2-165 grown at three different initial pH values (5.5, 6.2, 6.7) (Kettle et
584 al 2015 and Sylvia Duncan, personal communication).

585

586 **Table 1.** Capabilities for butyrate and propionate production among dominant bacterial species detected in
587 faecal samples of human subjects (Qin *et al.*, 2010; Zhernakova *et al.*, 2016)

Phylum (family)	species	Butyrate ¹	Propionate ²
Bacteroidetes (Bacteroidaceae)	<i>Bacteroides uniformis</i>	-	+ (Suc)
	<i>Bacteroides vulgatus</i>	-	+ (Suc)
Bacteroidetes (Prevotellaceae)	<i>Prevotella copri</i>	-	+ (Suc)
Bacteroidetes (Rikenellaceae)	<i>Alistipes putredinis</i>	-	+ (Suc)
Firmicutes (Lachnospiraceae)	<i>Eubacterium rectale</i>	+ (CoAT)	-
	<i>Roseburia inulinivorans</i>	+ (CoAT)	+ (Pdu)
	<i>Roseburia intestinalis</i>	+ (CoAT)	-
	<i>Dorea longicatena</i>	-	-
	<i>Eubacterium hallii</i>	+ (CoAT)	+ (Pdu)
	<i>Anaerostipes hadrus</i>	+ (CoAT)	-
	<i>Ruminococcus torques</i>	-	-
	<i>Coprococcus eutactus</i>	+ (ButK)	-
	<i>Blautia obeum</i>	-	+ (Pdu)
	<i>Dorea formicigenerans</i>	-	-
	<i>Coprococcus catus</i>	+ (CoAT)	+ (Acr)
	<i>Faecalibacterium prausnitzii</i>	+ (CoAT)	-
	<i>Subdoligranulum variabile</i>	+ (ButK)	-
Firmicutes (Ruminococcaceae)	<i>Ruminococcus bromii</i>	-	-
	<i>Eubacterium siraeum</i>	-	-
Firmicutes (Veillonellaceae)	<i>Dialister invisus</i>	-	+ (Suc)
Firmicutes (Acidaminococcaceae)	<i>Phascolarctobacterium succinatutens</i>	-	+ (Suc)
Firmicutes (Erysipelotrichaceae)	<i>Eubacterium biforme</i> ³	+ (CoAT)	-
Actinobacteria (Bifidobacteriaceae)	<i>Bifidobacterium adolescentis</i>	-	-

	<i>Bifidobacterium longum</i>	-	-
Actinobacteria (Coriobacteriaceae)	<i>Collinsella aerofaciens</i>	-	-
Verrucomicrobia (Verrucomicrobiaceae)	<i>Akkermansia muciniphila</i>	-	+ (Suc)

588

589 ¹-, absent; +, present; ButK, butyrate kinase route; CoAT, butyryl-CoA:acetate CoA-transferase route.

590 ²-, absent; +, present; Acr, acrylate pathway; Pdu, 1,2-propanediol pathway; Suc, succinate pathway

591 (succinate may be the major product formed instead of propionate in some species and/or under some

592 growth conditions).

593 ³Reclassified as *Holdemanella bififormis* (De Maesschalck *et al.*, 2014). CoA-transferase route is proposed

594 based on closely related butyrate producers within the Erysipelotrichaceae (see also main text, section on

595 CoA-transferases).

596

597

598 **Table 2.** Activity of the butyryl-CoA:acetate CoA-transferase of *Roseburia hominis* A2-183 (purified
 599 recombinant *butCoAT* gene product expressed in *Escherichia coli* (Charrier *et al.*, 2006)).

600

	K_m [mM]	V_{max} [μ mol/min/mg protein]	Inhibition by competition with acetate [%] ¹
acetate	6.4		
butyryl CoA	0.098	112	
propionyl CoA	0.099	51	
Butyrate			75
Propionate			70
Isobutyrate			56
Valerate			28

601 ¹No significant inhibition was found for caproate, 3-hydroxybutyrate, 4-hydroxybutyrate, 4-aminobutyrate,
 602 lactate, acetoacetate and succinate.

603

Fig. 1




Fig. 2





Fig. 3



606

607

Fig. 4

